

U.S. Patent Application

5

10

Genomeceutical and/or Enzymatic Composition
and Method for Treating Autism

15

Inventor(s): Mark A. Brudnak

0007615-050101
FOI090-47922860

Genomeceutical and/or Enzymatic Composition
and Method for Treating Autism

5

Inventor(s): Mark A. Brudnak

CROSS REFERENCE TO RELATED APPLICATIONS

This application claims the benefit of U.S.
Provisional Application No. 60/208,800, filed June 1,
10 2000, and having the same inventor as above and entitled
Enzyme Formulation for Treating Autism.

FIELD OF THE INVENTION

The present invention relates to treating autism and,
15 more specifically, to treating autism with genomeceutical
based treatments. The present invention also includes the
use of phytase and like substances and formulations
containing one or more of genomeceutical, enzymatic and
phytase-like compounds for treating autism.

20

BACKGROUND OF THE INVENTION

Autism may be defined as a condition, usually present
from childhood, that is characterized by self-absorption,
a reduced ability to respond to or communicate with the
25 outside world and behavioral dysfunction. An autistic
individual may suffer from several maladies with the
accumulated symptoms being categorized as autism spectrum
disorders, referred to in the field as autism or ASD.
Symptoms of autism include stimming, reduced eye contact,
30 perseveration (repeating same activity for long periods),
poor communication and social skills and heightened sound
sensitivity, amongst others.

It has been estimated that from 1 in 2,000 to 1 in 300 persons suffer from autism with an initial manifestation of symptoms by age three. It is of interest to note that the overall percentage of persons exhibiting symptoms of autism is increasing, in some instances dramatically. As discussed below, this rise may be due in part to an increase in the percentage of persons receiving childhood vaccinations. Males are more likely to suffer from autism than females.

There are several theories related to the initial and continued cause(s) of autism. One theory relates to infection or inflammation in the stomach and/or intestines early in life and maladaptation of the immune system and other tissues to this inflammation.

Historically, the immune system has been thought to develop early in life through education of the cellular components within the thymus (T-cells) and spleen (B-cells). There, the various cells are trained to recognize self vs. non-self antigens and to take the appropriate action, or lack thereof, in response to any exposure. In the last decade, however, a new class of immune cells has been intensively studied. These cells never "see" the thymus or spleen and are educated solely in the intestinal tissues from which they arise. These intraepithelial lymphocytes (IEL), or "intestinal" epithelial lymphocytes, are an athymically derived T-cell subset expressing the $\gamma\delta$ TCR-CD3 complex along with CD8. The $\gamma\delta$ T cells in the epithelial tissues do not circulate as their $\alpha\beta$ T cell (thymically derived) relatives do.

Though the subject of much speculation, the IELs are thought to be a primordial immune system and the first to function in life. IELs have been shown to bind to mycobacterial antigens that are protease resistant and

09072615 050104
TOT090 07092615

appear to have been selected to respond quickly to unique immune challenges. Since the mycobacterial antigens are protease resistant proteins, immune cells responsible for their removal would not benefit from an increased
5 production of protease genes. The immune cells would tend to conserve energy and redirect efforts by, for example, turning off protease genes. Thus, various protease genes, including those that appear to be effective in breaking down pre-opioid compounds that are linked to autism
10 (discussed below), may be down-regulated. In fact, DPPIV has been shown to be down regulated in autistics and is currently being used as a diagnostic marker for the disease. In addition, IELs may encourage apoptosis or quiescence of intestinal cells. This cell death or
15 disablement may lead to the "leaky" intestinal wall condition often associated with autism and ASD. Down regulation of protease genes and formation of "leaky" intestines are examples of the apparently maladaptive response mentioned above.

20 It is believed that these or other conditions cause the digestive tract of a person with autism to function sub-optimally. Two important pioneers of this work, Reichelt and Shattock, observed a significant correlation between the symptoms of autism and an impaired ability to
25 adequately digest peptides/proteins from dairy (casein) and wheat (gluten). During digestion, pre-opioid type compounds in the diet, typically from casein and gluten, are thought to be activated due to an incomplete breakdown of proteins. These exorphins (i.e.,
30 casomorphins and gluteomorphins or gliadorphin) are then easily transferred across the lumen of the gut into the circulation where they exert opioid-type action on the brain.

09872615 030401
T07090 5792860

In theory, no enzyme digestion in the body is ever 100% complete. Statistically, some protein or peptide fragments will escape the digestive process and be absorbed. The larger fragments may be transferred across the lumen of the gut via the M-cells and active transport while the smaller fragments may simply diffuse. Coupling these phenomena with the characteristically "leaky" intestinal tract of an autistic, it becomes readily apparent how biologically significant quantities of peptides and, more specifically, exorphins can enter the circulation.

The exorphins are recalcitrant to endogenous proteolytic enzyme digestion due to the presence of a proline in the penultimate position of the peptide. The body relies on its own production of dipeptidyl-peptidase IV (DPPIV), in the gut and other cells, for the digestion of exorphins. DPPIV is a serine exo-peptidase that cleaves Xaa-Pro dipeptides from the N-terminus of oligo- and polypeptides. It was first reported as glycylproline naphthylamidase and has been named dipeptidyl aminopeptidase IV or postproline dipeptidyl peptidase IV in early work. As alluded to above, it is suspected that genes which produce DPPIV are down regulated in autistic individuals.

To compensate for the apparent lack of sufficient quantities of DPPIV and to generally rebuild proper functioning of an autistic individual's intestinal tract with regard to absorption and digestion, different approaches have been employed. Of these, enzyme therapy and probiotic supplementation have been favored and met a degree of success. Enzyme therapy has typically been based on supplementation with large amounts of proteases from the different categories of proteolytic enzymes and

these have included acid or carboxyl peptidases, peptidases with both endo- and exo-peptidase activity, and serine, cysteine and zinc proteases. More recently, exogenous DPPIV from animal (usually cow or pig) and plant sources has been utilized. While enzyme therapy has had limited success, it is disadvantageous, amongst other reasons, in that many proteases, including DPPIV, are broken down in the stomach and do not reach the intestines in a functional state.

Probiotic supplementation has focused mainly on trying to rebuild the intestinal wall via a restoration of the naturally occurring bacterial flora. Common intestinal microflora are isolated from the human gut and cultured to form a "probiotic culture." Two approaches to the treatment of autistic dysbiosis using probiotic cultures have emerged. The first uses a wide variety of organisms and is something of a "shot-gun" approach. The second employs a more targeted approach by supplying very large numbers of viable organisms as high as 100 billion/gram. Both approaches have been based on clinically observed and published therapeutic benefits of probiotics and while they have met with limited success relative to no probiotic treatment, they do not adequately treat the syndromes associated with autism, suggesting that exorphin digestion is more than just an issue of microflora composition.

A need thus exists for a more effective manner of breaking down exorphins in the stomach and the intestinal tract of an autistic individual. A need also exists to increase expression of DPPIV and DPPIV like compounds from within an autistic individual's own body, i.e., overcoming the apparent down regulation of these or related genes. A need further exists for increasing the

bioavailability of minerals that facilitate digestion reactions.

SUMMARY OF THE INVENTION

5 Accordingly, it is an object of the present invention to increase expression of DPPIV and/or related compounds in the gut of an individual with autism.

 It is another object of the present invention to utilize genomeceutical materials that modify expression of
10 DPPIV, QPP or another compound whose regulation is beneficial in treating autism.

 It is another object of the present invention to provide various formulations that include genomeceutical material and one or more proteases/peptidases for treating
15 autism.

 It is also an object of the present invention to provide various formulations for treating autism that include phytase (or phytase-like substances) and protease and/or genomeceutical material.

20 It is a further object of the present invention to provide various formulations for treating autism that include phospholipids, disaccharidases and/or lipases.

 These and related objects of the present invention are achieved by use of a genomeceutical composition and
25 method for treating autism as described herein.

 The attainment of the foregoing and related advantages and features of the invention should be more readily apparent to those skilled in the art, after review of the following more detailed description of the
30 invention taken together with the drawings.

09373615-050101
T01090-01922860

10

15

20

30

increase expression of DPPIV in human intestinal wall cells and elsewhere. The increased amount of DPPIV would in turn achieve sufficient digestion of the pre-opioid exorphins to significantly reduce the symptoms of the autistic individual ingesting the galactose based composition.

In addition to galactose, the present invention may also include the use of other GCs, various proteases and/or peptidases, compounds that increase up take of substances that facilitate gene expression, phospholipids, disacchradases, lipases, related compounds and combinations of these items. Various representative, but not exclusive, embodiments of the present invention are now discussed in more detail.

Formulation A

In one embodiment of the present invention, termed formulation A for purposes of the present discussion, galactose is provided as a genomeceutical to increase expression of DPPIV. The galactose is preferably combined with other substances to enhance overall effectiveness. The other substances in the preferred embodiment of formulation A include: acid fast protease (AFP); bromelain and/or papain; peptidase concentrate; lactase and phytase.

If, in autistic individuals, the DPPIV gene has been silenced or attenuated (i.e., down-regulated), then the addition of galactose has the potential to reverse or circumvent the down-regulation. This positive regulation may occur not only in human intestinal wall cells (enterocytes), but in other cells where DPPIV or DPPIV-like enzymes are expressed (where a DPPIV-like enzyme is an enzyme that cleaves proline-containing peptide bonds in exorphins). It should also be recognized that DPPIV or

like expression in cells other than human enterocyte cells may proved to be as or more beneficial than enterocyte cell expression. These other cells include, but not limited to, peripheral blood (immune) cells and
5 other cells with suitable surface architectures and signaling cascades.

AFP is representative of a class of proteases that have high acid stability and function in the stomach to hydrolyze large proteins into smaller peptides. The
10 presence of a large amount of this enzyme is important in promoting rapid hydrolyzation of large proteins. The preferred result is that exorphins are formed early (in the stomach) and are digested (also in the stomach) before they reach a point of absorption (in the
15 intestines). Driving the reaction forward with large amounts of acid stable enzyme in the stomach allows more time for the body's endogenous enzymes, that have evolved to specifically digest exorphins, to work.

To further assist with this digestion a cystein
20 protease is preferably used. Bromelain and papain are examples of a cystein protease. Bromelain is preferred over papain because research has suggested that bromelain has a wider specificity and function than papain. It has also been demonstrated that bromelain is an effective
25 anti-inflammatory which may be significant in reducing the "leaky gut" characteristic of autistic individuals.

A peptidase concentrate component is preferably provided that exhibits endo- and exo-peptidase activity. It is further preferred that the peptidase concentrate
30 mimics DPPIV activity and hence provides further exorphin digestion. A suitable peptidase concentrate, amongst others, is the Case-Glutenase concentrate available

09072513060101

commercially from Kirkman Laboratories in Wilsonville,
OR.

Lactase is a disaccharidase that cleaves lactose into
its component sugars fructose and galactose. The
5 provision of lactase permits utilization of Formulation A
(or others herein) by lactose intolerant people and
increases that amount of available galactose. Cleaving
lactose and sucrose and other disaccharides also reduces
osmotic pressure in the intestines.

10 Phytase is preferably added for its ability to digest
phytic acid which is present in plants such as corn,
rice, wheat, soybean, and other beans, etc. Phytic acid
can negatively affect absorption of minerals such as
zinc, calcium, magnesium, copper, manganese, and iron.
15 Phytase supplementation results in greater
bioavailability of these minerals.

The components are generally available commercially
and are preferably provided in a dry form, mixed and
encapsulated, though other delivery methods may be
20 utilized without departing from the present invention.
The capsules are preferably taken with food. While the
presence and concentration of the above ingredients may
vary widely (as discussed to some extent herein below and
as recognized by one skilled in the art given the
25 teachings herein), in one embodiment formulation A may
contain the following:

- | | |
|-----------------------------|---------------|
| 1. galactose | 100mg; |
| 2. bromelain concentrate | 230 BTU; |
| 3. acid fast protease | 100 SAPU; |
| 30 4. peptidase concentrate | 10,000 AU; |
| 5. lactase | 300 LACU; and |
| 6. phytase | 125 PU. |

It should be recognized that these values are intended to be representative and in no way limit to the present invention.

This formulation A was given to 22 autism sufferers over a twelve-week period. Thirteen behavioral parameters were monitored including eye contact, socialization, attention, mood, hyperactivity, anxiety/compulsion, stimming, comprehension, speech, sound sensitivity, digestion, sleep and perseveration. Observers scored the patients every two-weeks for the twelve-week period. A Student's T-test was used for statistical significance determination.

Overwhelming positive trends were seen for each parameter with the greatest improvements in socialization and hyperactivity (90% and 80%, respectively) and the lowest improvements in stimming, speech and sound sensitivity, each scoring around 50% improvement.

The functioning of galactose is believed to be at at least two levels. A first level is as a genomeceutical where it is believed to be increasing the gut expression of the DPPIV gene. This increased expression allows for a greater level of the DPPIV enzyme in human enterocytes and other cells which in turn achieves a more thorough breakdown of any exorphins.

25 A second level is that galactose serves as a fuel
source of the beneficial microflora (i.e., probiotics) in
the gut. This is important because the probiotic
organisms themselves contain enzymes capable of
breaking-down the subject exorphins. Varmanen et al
30 recently showed that probiotic organisms, currently
utilized as health supplements, contain an analogue
of the DPPIV enzyme (e.g., PepX) which is known to be
able to digest exorphins. With over 10^{11}

microorganisms in the gut, the contribution of probiotic enzymatic activity may exceed that of the enterocytes. It is well documented that galactose is a prebiotic (i.e., stimulates growth of probiotics) and can increase the number of probiotics in the gut.

It is theorized thus that the level of self-produced DPPIV enzyme in the intestinal tract and the level of DPPIV-type activity (contributed by bacterial flora) are increased. It is further theorized that the ability of the AFP to cause early exorphin formation (in the stomach) and the increased levels of DPPIV production result in sufficient exorphin digestion in the gut that the levels of exorphin absorption drop below a threshold required for manifestation of the noted autistic symptoms.

Variations of Formulation A

It should be recognized that while Formulation A preferably has at least the listed six ingredients, the present invention may include combinations of less than all of the listed ingredients, as determined by the limits of prior art. For example, the combination of a GC for treating autism and proteases/peptidase is within the present invention as is the combination of a GC for treating autism and phytase or a phytase like compound. If a patient does not have milk in their diet or is not lactose intolerant then lactase may not be necessary. Further variations are suggested elsewhere herein and others yet would be apparent to skilled practitioners given the teachings herein. Given the exponential combination of ingredients all variations are not specifically called out, though it should be recognized

that they are intended to fall within the present invention.

Beta Glucans and QPP

5 Recently, a new peptidase (quiescent cell proline dipeptidase or QPP) from peripheral blood mononuclear cells (PBMCs) has been cloned and investigated. QPP is structurally and functionally related to DPPIV, but is more often found in the peripheral blood cells.
10 Interestingly, QPP shares little sequence homology with DPPIV. These enzymes (DPPIV, QPP and others) are among the few enzymes that have the ability to cleave proline-containing peptide bonds. The exorphin opioid type peptides containing proline are generally resistant to
15 degradation as smaller peptides.

Notwithstanding increased levels of DPPIV produced by the enterocytes by the addition of galactose, there may still be opioid peptides that reach circulation. If the QPP gene can be up-regulated, then exorphins that do
20 manage to survive across the lumen of the gut are more likely to be digested once inside the circulatory system.

At the present time, there is preliminary data that suggests that stimulation of the TCR causes an up-regulation of the QPP gene. The TCR can be stimulated by
25 a number of known factors. These include, but are not limited to, antibodies, lectins, and macrophages. While certain natural and potentially safe lectins may exist and be shown useful in the future, current technology and regulatory issues makes mammalian lectins impractical.
30 Also, as shown by empirical data, plant lectins can be extremely dangerous. Antibodies offer some potential, but there are also many problems associated with their use. These range from their ethical production to safety.

Encouragement of macrophage participation in accordance with the present invention solves most of these problems and presents an opportunity for therapeutic application of genomeceuticals.

5 It is well known that in the gut, cellular components of bacteria, fungi, and molds can stimulate gut-associated macrophages. These macrophages constantly sample and survey the gastrointestinal tract for foreign antigens. The cellular components responsible for the
10 stimulation of macrophages have been well documented and include not just proteins, but carbohydrates such as the beta glucans present in cell walls of microorganisms.

When macrophages are exposed to beta glucans, such as the yeast cell wall beta-1,3/1,6-glucan, they become
15 non-specifically stimulated so that subsequent antigen exposure results in a much more robust immune response than would otherwise happen. Among those responses is the secretion of cytokines and the processing and subsequent presentation of antigens to T-cells. The role of antigen
20 presentation is accomplished through interaction with the TCR. The bound TCR then transduces a signal to the nucleus directing an altered expression pattern which includes modified expression of available QPP. QPP is then secreted into circulation and is free to digest any
25 exorphins present. Glucans have also been shown to up-regulate transcription factors in the cell nucleus. Hence, glucans can serve as a genomeceutical for increasing expression of QPP.

Accordingly, in another embodiment, the present
30 invention includes one or more of the components of Formulation A with glucans and/or QPP (or their analogues).

0087261E 060104
TOT090 4922860

Other Genomeceuticals/Sugars

Lactose, fructose, sucrose, glucose, etc., are sugars that may be present in milk, as is galactose. The presence of one or more of these sugars may cause the intestinal
5 cells to produce enzymes, including DPPIV like enzymes, that are effective in breaking down milk proteins and milk protein by-products (such as exorphins, amongst others). Thus, while galactose is a preferred sugar, it is possible that one or more of these other sugars and/or related
10 compounds may exhibit efficiency in promoting expression of DPPIV like enzymes. Furthermore, the provision of lactose and lactase would provide galactose as a by-product.

Thus, including one of these sugars or a related
15 compound as a substitute for or in addition to the galactose of Formulation A or with any variation of the components or Formulation A or their analogues or any other variation alluded to herein is within the present invention.

20

Protease/Peptidase

While Formulation A includes a particular protease and peptidase arrangement, it should be recognized that other protease and peptidase arrangements may be utilized
25 without departing from the present invention. These include carboxyl, serine, and zinc proteases and any other proteases that would be efficacious and suitable. Other peptidase concentrates or specific peptidase compounds may be utilized.

30

These other proteases and peptidase may be used as substitutes for or in addition to the component(s) of Formulation A or other components mentioned herein.

Phytase

In addition Formulation A, the present invention includes the combination of phytase and a genomeceutical for treating autism and the combination of phytase and a
5 protease and/or peptidase for treating autism.

Various phytase like compounds are suitable for use in the present invention, where phytase like refers to compounds that increase the bioavailability of one or more of the listed minerals.

10

Phospholipids

Phospholipids may be provided in combination with less, all or more of the ingredients of Formulation A or other variations alluded to herein. Phospholipids would
15 serve to rebuild cell walls and help heal the leaky guts of autistic individuals. A representative dose of phospholipids could be 300 mg of phosphotidyl serine or the like.

Disaccharidase

Lactase, sucrase and/or other disaccharidases may be provided in accordance with the present invention. The functions of these compounds is discussed in part above with the discussion of lactase. The disaccharidase(s) may
25 be provided as a substitute for or in addition to other components and other varied formulations discussed or alluded to herein.

Lipase and Other Enzymes

30 Lipase, amylase and other related enzymes may also be provided in accordance with the present invention. Amylase liberates glucose from carbohydrates, and the liberated carbohydrate may serve as a genomeceutical as discussed

09872615 060101
TOT090 57924860

Lipase serves to digest fat. Undigested fats lead to loose stools which are a condition associated with autism.

While the invention has been described in connection with specific embodiments thereof, it will be understood that it is capable of further modification, and this application is intended to cover any variations, uses, or adaptations of the invention following, in general, the principles of the invention and including such departures from the present disclosure as come within known or customary practice in the art to which the invention pertains and as may be applied to the essential features hereinbefore set forth, and as fall within the scope of the invention and the limits of the appended claims.